

Density dependence at the time of spawning: disentangling density-dependent effects in the life histories of fish

Akinori Takasuka¹, Michio Yoneda², Yoshioki Oozeki³

Japan Fisheries Research and Education Agency

¹National Research Institute of Fisheries Science

²National Research Institute of Fisheries and Environment of Inland Sea

³Headquarters


PICES 2018 Annual Meeting

Yokohama, Japan

Session 1, October 29, 2018

Spawner–recruitment

Spawner–recruitment relationship of the Pacific stock of **Japanese sardine** (Yukami *et al.* 2017 Stock assessment report)



Backgrounds

- The asymptotic pattern represents density-dependent effects on recruitment (i.e. self-regulating processes).
- Any variability from the asymptotic pattern represents density-independent effects on recruitment (i.e. environmentally-regulated processes).
- Evidence of density-dependent effects on juvenile growth and maturation.

Interpretation

- Density-dependent mortality from hatching to recruitment.
- Density-independent mortality due to climate impacts.

Paradigm

A paradigm of fisheries science

“Spawner biomass is a suitable proxy for reproductive potential of a fish stock.”

Leggett & Frank (2008) “Paradigms in fisheries oceanography”
Oceanography and Marine Biology: An Annual Review



“SSB–TEP proportionality” paradigm


“Spawning stock biomass (SSB) and total egg production (TEP) are proportional to each other.”

A basic premise underlying the spawner–recruitment models for fisheries management and studies on recruitment mechanisms of fish.

Ricker (1954), Beverton & Holt (1957), Barrowman & Myers (2000)

Spawner–recruitment

Spawner–recruitment relationship of the Pacific stock of **Japanese sardine** (Yukami *et al.* 2017 Stock assessment report)



Backgrounds

- The asymptotic pattern represents density-dependent effects on recruitment (i.e. self-regulating processes).
- Any variability from the asymptotic pattern represents density-independent effects on recruitment (i.e. environmentally-regulated processes).
- Evidence of density-dependent effects on juvenile growth and maturation.

Reproductive potential of a stock



Total egg production (TEP)



Spawning stock biomass (SSB)

Interpretation

- Density-dependent mortality from hatching to recruitment.
- Density-independent mortality due to climate impacts.

Paradigm





“SSB–TEP proportionality” paradigm

- Progress in studies on maternal effects on reproductive potential has led to doubt about the paradigm.
“Bigger, older, fatter, and repeat spawners produce larger and more eggs”
- Nonetheless, a direct test of the paradigm at multidecadal scales has been difficult because of data limitations in the stock assessment systems worldwide.
- It is difficult to prepare TEP data independent of SSB data.

In the present study, ...

- We directly tested the paradigm based on a combination of two independent time series (SSB and TEP), using sardine and anchovy as model species.
- To show that the “SSB–TEP proportionality” is distorted by density dependence in egg production.

Egg surveys





Egg surveys covering the spawning grounds during the high- and low-biomass periods of sardine and anchovy in the western North Pacific from 1978 to 2015 (38 years). The data set includes a total of 145,157 vertical tows of plankton nets.

Egg production

Annual total egg production (TEP)

1. Monthly mean egg density for each 15' × 15' (latitude × longitude) square
2. Monthly egg abundance for each square = monthly mean egg density for each square × square area
3. The parameters considered to realistically assess the quantitative data:
 - Egg developmental stage
 - Egg incubation time (a function of temperature)
 - Egg survival rate
4. Monthly egg abundance = sum of monthly egg abundance for all the squares
5. Annual total egg production (TEP) = sum of monthly egg abundance for all the months
 - **Sardine**: October of the previous year to September of the year
 - **Anchovy**: January to December of the year




Examples of distributions of monthly egg production.


Biomass and TEP

A combination of two independent 38-year time series

Fishery-dependent stock assessment data
(derived from stock assessment reports)



Fishery-independent egg survey data




Multidecadal changes in biomass and annual total egg production (TEP) of the Pacific stocks of sardine and anchovy.

TEP vs SSB


Direct test of the “SSB–TEP proportionality” paradigm based on a combination of two independent 38-year time series

- Spawning stock biomass (SSB): fishery-dependent stock assessment data
- Annual total egg production (TEP): fishery-independent egg survey data



$$y = 114 \times 10^6 x$$

($n = 37$, $R^2 = 0.636$, $p < 0.001$)



$$y = 7792 \times 10^6 x$$

($n = 38$, $R^2 = 0.917$, $p < 0.001$)

Relationships of annual total egg production (TEP) to spawning stock biomass (SSB) for sardine and anchovy.


TEPPS

Annual total egg production per spawner individual (TEPPS)

= Annual total egg production (TEP) / spawning stock abundance (SSA)


= How many eggs are produced from an individual spawner for a year

= Annual realized fecundity (integrating batch fecundity, spawning fraction, etc.)



$$y = 7,400 \times x$$

($n = 37$, $R^2 = 0.353$, $p < 0.001$)







$$y = 111,605 \times x$$

($n = 38$, $R^2 = 0.754$, $p < 0.001$)

Relationships of annual total egg production (TEP) to spawning stock abundance (SSA) for **sardine** and **anchovy**.


TEPPS vs SSB



Relationships of annual total egg production per spawner (TEPPS) to spawning stock biomass (SSB) for **sardine** and **anchovy**.


TEPPS vs SSB

- A significantly negative relationship appeared when TEPPS of anchovy was related to SSB of sardine.
- Hypothesis: TEPPS of anchovy is influenced by sardine biomass.
- Analysis: seasonal/regional TEPPS of anchovy versus SSB of sardine.




Relationships of annual total egg production per spawner (TEPPS) of anchovy to spawning stock biomass (SSB) of sardine.

Seasons



Examples of distributions of monthly egg production of sardine and anchovy off the Pacific coast of Japan.

Regions




The six regions were classified by the location of the Kuroshio axis. The regions spatially and temporally change with the spatial (mesoscale) and temporal (daily) variability of the Kuroshio axis.

Definitions

- IK: Inshore side of the Kuroshio axis region
- KA: The Kuroshio axis region
- OK: Offshore side of the Kuroshio axis region
- KE: The Kuroshio Extension region
- KES: The Kuroshio Extension southern region
- KOT: The Kuroshio–Oyashio transition region

Examples of regions classified by the location of the Kuroshio axis.

TEPPS vs SSB



Relationships of seasonal/regional total egg production per spawner (TEPPS) of anchovy to spawning stock biomass (SSB) of sardine.


TEPPSW

Annual total egg production per spawner unit weight (TEPPSW)

= Annual total egg production (TEP) / spawning stock biomass (SSB)


= How many eggs are produced from a body mass unit weight of spawner for a year

= Annual relative fecundity (annual realized fecundity relative to body mass of spawner)



$$y = 114 \times 10^6 x$$

($n = 37$, $R^2 = 0.636$, $p < 0.001$)







$$y = 7792 \times 10^6 x$$

($n = 38$, $R^2 = 0.917$, $p < 0.001$)

Relationships of annual total egg production (TEP) to spawning stock biomass (SSB) for **sardine** and **anchovy**.


TEPPSW vs SSB



Relationships of annual total egg production per spawner unit weight (TEPPSW) to spawning stock biomass (SSB) for **sardine** and **anchovy**.


TEPPSW vs SSB

- No significant relationship appeared when TEPPS of anchovy was related to SSB of sardine.
- However, negative relationships of TEPPS of anchovy to SSB of sardine were confirmed by an analysis on seasonal/regional TEPPS of anchovy versus SSB of sardine.



Relationships of annual total egg production per spawner unit weight (TEPPSW) of anchovy to spawning stock biomass (SSB) of sardine.

TEPPSW vs SSB



Relationships of seasonal/regional total egg production per spawner unit weight (TEPPSW) of anchovy to spawning stock biomass (SSB) of sardine.

Summary I

Direct test of the paradigm

- We directly tested the “SSB–TEP proportionality” paradigm based on a combination of two independent time series.
- The “SSB–TEP proportionality” is distorted by intraspecific and interspecific density dependence in TEPPS/TEPPSW (egg production).

Density dependence in TEPPS/TEPPSW


- The operation markedly differed between the species.
Sardine: intraspecific (self-density-dependent)
Anchovy: interspecific (**sardine**-density-dependent)
- The phenomena would be attributable to maternal effects through intraspecific and interspecific competition for food resources required as energy for reproduction.
- Density dependence in egg production was previously unaccounted for in spawner–recruitment relationships.
- How does the density dependence in egg production change the current understanding of density-dependent processes in the life history of fish?

Summary I

- The phenomena would be attributable to maternal effects through intraspecific and interspecific competition for food resources required as energy for reproduction.


(1) Trophic overlap

Sardine and **anchovy** have strong trophic overlaps throughout their life histories.



(2) Population fluctuation

The maximum biomass is 13 times greater for **sardine** (19.5×10^6 t) than for **anchovy** (1.5×10^6 t); the extent of population fluctuation is 20 times greater for **sardine** (225) than for **anchovy** (11).



(3) Energy allocation strategy for reproduction


Sardine: “capital–income” breeder (expending stored energy reserves)

Anchovy: “income” breeder (expending energy recently acquired)



Disentangling


Disentangling density-dependent effects on egg production and survival from egg to recruitment



Relationships among spawning stock biomass (SSB), annual total egg production (TEP), and recruitment (R) for **sardine** and **anchovy**.

Disentangling


Disentangling density-dependent effects on egg production and survival from egg to recruitment



Relationships of recruitment per spawning stock abundance (RPS), annual total egg production per spawner (TEPPS), and recruitment per total egg production (RPE) to spawning stock biomass (SSB) and total egg production (TEP).

Disentangling

Disentangling density-dependent effects on egg production and survival from egg to recruitment



Relationships of recruitment per spawning stock abundance (RPS), annual total egg production per spawner (TEPPS), and recruitment per total egg production (RPE) to spawning stock biomass (SSB) and total egg production (TEP).

Summary II

Disentangling density-dependent effects

- The existence of density dependence in egg production could change the current understanding of density-dependent processes in the life history of fish.
- Proposing to consider a shift from SSB-based to TEP-based strategies in fisheries management and recruitment studies.

1. Implications for fisheries management

- Determining biological reference points based on TEP–R relationships rather than SSB–R relationships.
- Considering interspecific interactions in biological reference points.

2. Applications to recruitment studies

- Using TEP-based survival indices for the process from hatching to recruitment.

3. Contributions to numerical modelling


- Closing the life cycle by quantifying the SSB–TEP relationships.
- Constructing multispecies or interactive models.

Future perspectives

- Application to other pelagic and demersal species.
- Promoting and implementing fishery-independent egg surveys.

Received: 28 March 2018 | Revised: 21 August 2018 | Accepted: 3 September 2018

DOI: 10.1111/faf.12327

ORIGINAL ARTICLEWILEY 

Density dependence in total egg production per spawner for marine fish

Akinori Takasuka¹  | Michio Yoneda² | Yoshioki Oozeki³

¹National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency, Yokohama, Japan

²National Research Institute of Fisheries and Environment of Inland Sea, Japan Fisheries Research and Education Agency, Imabari, Japan

³Japan Fisheries Research and Education Agency, Yokohama, Japan

Correspondence

Akinori Takasuka, National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency, Yokohama, Japan

Abstract

A paradigm of fisheries science holds that spawning stock biomass (SSB) is directly proportional to total egg production (TEP) of fish stocks. This “SSB-TEP proportionality” paradigm has been a basic premise underlying the spawner–recruitment models for fisheries management and numerous studies on recruitment mechanisms of fish. Studies on maternal effects on reproductive potential of a stock have progressed during the last few decades, leading to doubt concerning the paradigm. Nonetheless, a direct test of the paradigm at multidecadal scales has been difficult because of data limitations in the stock assessment systems worldwide. Here, we tested the paradigm for marine fish based on a novel combination of two independent 38-year time

Takasuka, A., Yoneda, M., Oozeki, Y. (2018) Density dependence in total egg production per spawner for marine fish. *Fish and Fisheries*.