How will climate driven shifts in coral reef communities influence their suitability as fish habitat?

Preliminary results of my 2nd PhD chapter

Ravi Maharaj

UBC, Institute for the Oceans and Fisheries
William Cheung, Daniel Pauly, Simon Donner and Rashid Sumaila





Coral reefs are important, but in danger

 Corals are important habitat for fish communities

 Coral fisheries support important local and global supply chains

 Sensitive to changes in their physical environment








Morphology-specific responses to thermal stress related to 'mass-transfer'

"...the potential exchange rate of gases and metabolites across corals' boundary layers."

- This rate is lower on larger, branching colonies
- Metabolic processes more vulnerable under environmental stress

van Woesik et al. 2012




Coral mortality leads to impacts on fish communities and fisheries

 Dead coral susceptible to physical and chemical erosion

Results in declining habitat complexity

Reduced complexity affects fish communities




MARINE ECOLOGY PROGRESS SERIES Mar Ecol Prog Ser

Published March 12

Regional variability in the sensitivity of Caribbean reef fish assemblages to ocean warming

Ravi R. Maharaj^{1,*}, Vicky W. Y. Lam¹, Daniel Pauly², William W. L. Cheung¹

¹Changing Ocean Research Unit, ²Sea Around Us, Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, BC V6T 1Z4, Canada




Research questions

1. How will projected changes in the oceanic environment influence shifts in the distribution of morphology groups?

1. How may this influence the quality of habitat provided to fish by reef communities?

Area of study – Caribbean Large Marine Ecosystem

- Not as speciose as Pacific
- Habitat quality declined significantly 1980-2000
- North: The Bahamas, US East,
 Cuba
- Central: Mexico, Honduras, Nicaragua, Belize
- South: Trinidad & Tobago,
 Venezuela, Colombia



Predict distribution of corals using Species Distribution Models


"...SDMs create representations of species' fundamental niche which can be extrapolated into space and time (Habitat Suitability Index)..."


- Species occurrence data
 - 63 species (56% of species with occurrence data)
- Environmental variables
 - 7 climatologies (1970-2000)
 - 2 climate scenarios (RCP 2.6 & 8.5)
 - 2 future periods (2050, 2100)
 - 3 earth system models (IPSL, MPI, GFDL)
- Used to model species, functional groups and habitat types
 - ANN, Maxent, GBM

Morphological categories and ranks

 Categories from the coral trait database (<u>www.coraltrait.org</u>)


- Morphology classes
 - Acroporid corals 3
 - Sub-massive and Massive corals 2
 - Non-acroporid branching and encrusting corals – 1 (Edinger and Risk, 2000)







Habitat Quality Index

Higher as HQ -> 6 Lower as HQ -> 0





Habitat Quality Regional Baseline (1970-2000)






Habitat Quality Future Projections







Area Trends



% Change in Mean Regional Habitat Quality



% Change by Area from Regional Baseline



Key observations

- RCP scenarios produce vastly different futures for habitat quality
- Stabilization of HQ under 2.6, but drastic declines under 8.5
- Patterns among areas are similar, but vary greatly in ΔMHQ

Preliminary conclusions

- Habitat management may be useful; spatial trends must be considered
- Central America may act as a climate refuge
- Other important variables may affect these trends
- Higher emissions scenario may require adaptation of local fisheries if fish communities are affected (more on that later next year...)

Future Analyses

- What are the shifts in coral communities by morphology and species?
- What are the key environmental variables?
- What are the consequences for fish communities?

Acknowledgements























