



Quantifying *Calanus finmarchicus* growth and dormancy using AARS activity


Lidia Yebra, Andrew G. Hirst, Santiago Hernández-León



Protein synthesis






Aminoacyl-tRNA synthetases mechanism



AARS assay

PPi reagent

Zooplankton homogenate \Rightarrow Decrease in absorbance at 340 nm



0

10 min


Advantages

- AARS activity can be assessed without incubating organisms
- Growth can be estimated between and within stages, moulting is not required
- The AARS method has been successfully calibrated as growth index for *Calanus helgolandicus* copepodites and adults
(Yebra et al., 2005, Mar. Biol.)

Calanus finmarchicus



Jean Francois St-Pierre




Growth and AARS activity


Location and differential characteristics of each experiment set.

	Espeland Station	St Lawrence	Irminger Sea	Labrador Sea
Sampling location	60°24'N, 05°18' E	48° N, 66° W	58-65° N, 22-42° W	42-44° N, 60-65° W
Experiment location	Espeland Laboratory	IML Laboratory	RRV Discovery	CCGS Hudson
Date	May 1998	August 2000	August 2002	May 2004
Stages	CIV-CVI	CV	CIV-CV	CI-CIII
Cultures T°C	8	8	5, 9	0-4
Food type	<i>Rhodomonas baltica</i>	<i>Rhodomonas baltica</i>	Natural food	Natural food
Food concentration	100-800 µgC/L	0-800 µgC/L	Field conc.	Field conc.


Growth and AARS activity



Growth and AARS activity




Dormancy and AARS activity



Dormancy and AARS activity

Calanus finmarchicus



Summary

- The AARS method is a suitable index of *Calanus finmarchicus* somatic growth across the North Atlantic ocean
- AARS activity can be successfully assayed on overwintering organisms
- The AARS method opens the possibility to study protein metabolism of deep sea *Calanus* populations

Special thanks to:

S. Alvarado

U. Båmstedt

G.-G. Chang

L. Harris

R.P. Harris

S. Hay

E. Head

S. Helland

P. Joly

F. Maps

J. Runge

Discovery & Hudson Crews

More info:

Yebra, Hirst and Hernández-León (2006) J. Plankton Res. 28(12): 1191-1198